

## Zooplankton diversity of Keenjhari Lake district Thatta

# Lashari K.H.<sup>1</sup>; Palh Z.A.<sup>1\*</sup>; Shaikh A.A.<sup>2</sup>; Bughio B.A.<sup>3</sup>; KhemtioD.<sup>3</sup>; Mashori D.B.<sup>4</sup>; Suhag R.<sup>1</sup>

Received: February 2024

Accepted: May 2024

#### Abstract

The goal of this study was to learn more about the zooplankton diversity in Keenjhar Lake. Keenjhar Lake is a tropical, man-made lake. It is around 120 kilometers, Outside of Karachi, at a distance of 24°47' N latitude and 68°02' E longitude (Ayoade and Aderogba, 2020). "Keenjhar" and "Sunheri," or "Green" and "Golden," respectively, two natural lakes were joined to create the lake. By blowing up the dividing hills in 1958, Keenjhar and Sunheri, two tiny depressions, were combined to create the "Kalri Lake.". Zooplanktons are significant fauna in water bodies that operate as pollution bioindicators and play a direct part in the food chain. There were a total of 9 different zooplankton species found. There were 06 Rotifera species, 01 Copepoda species, and 02 Cladocera species discovered among them. Brachionus calyciflourus (Pallas, 1834), Brachionus bidentata (Ahangar and Farooq, 2012), Lecane leontina (Aziz1 et al., 2014), Trichotria tetractis (Ehrenberg, 1830), and Trichocerca pusilla (Ehrenberg, 1830) were among the rotifers found (Jennings, 1903). Moina micrura (Lashari et al., 2021) was found in Copepoda, and Ceriodaphnia quadrangular (Müller, 1785) and Diaphanosoma birgei Ascomorpha saltans (Lashari et al., 2021), were found in Cladocera (Korineck 1981). Rotifera species are the most common zooplankton species, indicating that the lake water is contaminated.

Keywords: Rotifera, Copepoda, Cladocera, Diversity, Zooplankton and Thatta

<sup>1-</sup>Department of Fisheries and Aquatic Sciences, University of Sindh Jamshoro, Pakistan

<sup>2-</sup>Department of Chemistry Government Collage University Hyderabad, Pakistan

<sup>3-</sup>Department of Zoology University of Sindh Jamshoro, Pakistan

<sup>4-</sup>Department of Zoology Government Collage University Hyderabad, Pakistan

<sup>\*</sup>Corresponding author's Email: alizameer735@gmailcom

## Introduction

Zooplankton is a vital component of aquatic wildlife and a key component of the aquatic food chain. It also ensures that the aquatic ecosystem's biotic and abiotic components are in perfect balance. Freshwater zooplankton is made up of three primary kinds of Rotifers, Copepods, and Cladocera's are invertebrate creatures. They are widespread they play a critical part in the transfer of energy in all varieties of aquatic ecosystems. and the biomarker of pollutants. It has a middle place in the food chain, with many of them feeding on bacteria and algae, which are then eaten by a variety of invertebrates, fish, and birds (Zhang Honggang and Cui Baoshan). The variety of Understanding zooplankton and their ecology greatly aids in our comprehension of the fundamental makeup and overall functioning of aquatic environments. The population of zooplankton in a water body is also influenced by physical and chemical variables. The zooplanktons of several fresh water sources were studied by a number of researchers (Lashari et al., 2021) Investigated the variety of zooplankton in the river Kayadhu in Hingoli, Maharashtra. Rotifer Checklist was made by Lashari et al. (2014) after researching the population of rotifers in the Washim region. Pawar and Dabhade investigated the qualitative diversity of the rotifer population in the freshwater Katepurna reservoir in the district of Akola, Maharashtra, India (Lashari et al., 2008). Zooplankton diversity is a useful indication of changes in water

resources since it reflects the quality of the water. 7. (Lashari et al., 2008) conducted study in Washim. to examine the Maharashtra, India, zooplanktons of the freshwater environment. Because daily pollution is rising and has a direct influence on aquatic life, aquatic life biodiversity protection is a critical issue. Pollution and human activities have put biodiversity at jeopardy in recent years. Because biodiversity conservation is critical, it is necessary to maintain current knowledge on all aquatic species variety (Sunder et al., 2018). Because planktons are the primary source of food for economically significant fishes as well as recreational fishes the pace of fish stocking is determined by the density of planktons in a body of water.

## Materials and methods

Zooplankton diversity is a useful indication of changes in water resources since it reflects the quality of the water. Korai et al. (2008) conducted study in Washim, Maharashtra, India, to examine the zooplanktons of the freshwater environment. Because daily pollution is rising and has a direct influence on aquatic life, aquatic life biodiversity protection is a critical issue. Pollution activities and human have put biodiversity at jeopardy in recent years. Because biodiversity conservation is critical, it is necessary to maintain current knowledge on all aquatic species variety.

The concentrated zooplankton samples were moved to a new container with care. 2 to 3 drops of glycerin were applied to 5 ml of 4 percent formalin. To prevent zooplankton aggregation, a sprinkle of detergent powder was applied. The samples were gathered in separate glass phials with labels that included the name of the location, the date of sampling, the time of sampling, and other information. The researchers used a compound microscope to identify the zooplankton. Sorting and counting also done with a dissecting are microscope. Glass slides were used to hold the specimens while they were examined at magnifications ranging from 25 to 100X. Williamson (Korai et al., 2010), Dodson and Frey (Lashari, 2002) Tonapi (Lashari et al., 2008), Dodson and Frey (1991), APHA (1985) and Battish's systematic key were also employed, along with conventional identification methods and monographs. Population size was calculated using Lackey's drop count method from 1938.

#### **Results and discussion**

22 recorded zooplankton species identified from Keenjhar Lake (Lashari *et al.*, 2012) discovered 15 different species of Zooplankton. During the whole research period, Rotifer's population was dominating among familiar Zooplankton. The mean density of reported Protozoans ranged starting 2 org./L to 31 org./L at various sample locations. The largest density of protozoans was found in August, even as the lowest density was found in August (rainy season). Protozoan density varies seasonally the average value (Zaheer Khan et al., 2014). The graphic depicts the seasonal variation in Rotifer density. Throughout the investigation, the mean observed density of Cladocerans ranged from 4 to 84 org./L. In August, when they were at their densest, they were at their highest density. In May, when it was summer (rainy season). The graphic depicts the seasonal variation in Cladocera density. The reported average Copepod density over the study period was in the range of between 6 and 42 org./L. The month of May (summer) was this group's highest density, (Shahina and Azmi, 2019) whereas the month of August (rainy season) had its lowest density). Graphic depicts the seasonal variation in Copepoda density. Zooplankton density was found to be highest in the summer, intermediate in the winter, and lowest during the rainy season. Table 1 shows the seasonal average of reported Zooplankton density in Keenjhar Lake research area.

Protozoa	Genera	Species	Cladocera	Genera	Species
1	Arcella	Dentata caudatum	1	Alona	Rectangula
2	Paramecium	Dentata caudatum	2	Bosmina	Longirostris
			3	Ceriodaphnia	Reticulata
3	Vorticella	Dentata caudatum	4	Daphnia	Rarinata
			5	Moina	Brachiata
Copepoda	Genera	Species	Rotifera	Genera	Species
	Cyclops	Bicupidatus	1	Asplanchna	intermedia
	Macrocylops	Albidus	2	Asplanchna	intermedia

Zooplankton in Keenihar Lake that have been spotted on record.





Figure 4: Zooplankton density varies seasonally (Copepoda).

### Reference

- Adedolapo A. Ayoade and Ashiat Aderogba, 2020. Spatial and temporal distribution of plankton in a reservoir, southwestern tropical Nigeria. Egyptian Journal of Aquatic Biology Å *Fisheries* Zoology Department, Faculty of Science, Ain Shams University, Cairo, Egypt. ISSN 1110 - 6131, 24(5), 161 - 181.
- Ahangar, A. and Farooq, M., 2012. Zooplankton diversity with relation to trophic status in Anchar Lake, Kashmir. *International Journal of Current Research*, 4, 07, pp. 046-048.
- Farha Aziz1., Rafia Azmat and Farah Jabeen. 2014. Water quality and pollution burden of Keenjhar Lake, Thatta, Sindh, Pakistan. International Journal of Recent Scientific Research Research, 5, 4, pp.770-773.
- Khalid Hussain Lashari, S. Habhib Naqvi, Zameer Ali Palh. (2014). The effects of physiochemical parameters on planktonic species population of Keenjhar lake, district Thatta, Sindh, Pakistan. *American Journal of BioScience*, 2(2), 38-44. http://www.sciencepublishinggroup. com/j/ajbio.DOI:

10.11648/j.ajbio.20140202.13

- Khalid Hussain Lashari, Zameer Ali Palh, Shaharyar Brohi, Sujo Meghwar., (2021). Zooplankton variety in saline water: Investigation of different varieties of zooplankton in Keenjhar Lake of District Thatta. Univeristy of Sindh. *Journal of Animal Sciences*, 5(3), 31-34.
- Khalid Hussain Lashari, Zameer Ali
  Palh, Shaharyar Brohi, Sujo
  Meghwar., 2021. Assessment of water chemistry and phytoplankton: A case study of Keenjhar lake District Thatta, Sindh Sindh university. Research (Science Series), 53(3), 209-212.
- Korai, A.L., Lashari, K.H., Sahato,
  G.A. and Kazi, T.G., 2010.
  Histological lesions in gills of feral
  Cyprinids, related to uptake of
  waterborne toxicants from Keenjhar
  Lake. *Rev. Fisher. Sci.*, 18(2), 157-176.
- Korai, A.L., Sahato, G.A., Lashari,
  K.H. and Arbani S.N., 2008.
  Biodiversity in relation to physicochemical properties of Keenjhar Lake, Thatta district, Sindh, Pakistan. *Turk. J. Fish. Aquatic Sci.*, 8, 259-268.

- Lashari, K.H., 2002. Ecological studies of plankton of Keenjhar Lake, district Thatta, Sindh, Pakistan. M. Phil. dissertation, Department of Fresh Water Biology and Fisheries, University of Sindh, Jamshoro, Pakistan
- Lashari, K.H., Korai, A.L., Sahato, G.A. and Kazi, T.G., 2009. Limnological studies of Keenjhar Lake, District, Thatta, Sindh, Pakistan. *Pak. J. Anal. Environ. Chem.* 10(1-2), 39-47.
- Lashari, K.H., Sahato, G.A., Korai, A.L. and Kazi, T.G., 2008. Taxonomic study of chroocophyceae (Cyanophyta) in Keenjhar Lake, Distt: Thatta, Sindh, Pakistan. *Research Journal of Fisheries and Hydrobiology*, 3(1),11-21.
- Lashari, K.H., Sahato, G.A., Korai, A.L. and Kazi, T.G., 2008. Taxonomic study of Chroocophyceae (Cyanophyta) in Keenjhar Lake, Dist: Thatta, Sindh, Pakistan. *Res. J. Fisher. Hydrobiol.*, 3(1),11-21.
- Lashari, K.H., Sahato, G.A., Korai, A.L., S. Habib Naqvi, Palh, Z.A. and Urooj, N., 2012. Heavy metals

burden of Keenjhar Lake, District Thatta, Sindh, Pakistan. *Afri. J. Biotech.* 11(**59**), 12305-12313,

- Shahina Rao, B. Mukhtar and S.H. Hasany.2016. Abundance of freshwater zooplankton and Physicochemical parameters in Quaid-e-Azam Park Lake, Karachi, Pakistan. *Int. J. Biol. Biotech.*, 13 (2), 253-256.
- Shahina Rao; Azmi, M.A., 2019. Species composition and occurrence of Cladocera in Keenjhar lake, Sindh, Pakistan. *International Journal of Biology and Biotechnology*, 16, 2, pp.509-512.
- Shyam Sunder. Dr. Anand Kumar Khatri 2018. Zooplanktonic fauna in relation to physico-chemical properties in Ottu Reservoir, District Sirsa, Haryana, India. International Journal of Creative Research Thoughts (IJCRT), 6, 1.
- Zhang Honggang, Cui Baoshan., 2012. Species diversity and distribution for zooplankton in the intertidal wetlands of the Pearl River estuary, China. *Procedia Environmental Sciences*, 13, 2383 – 2393